

# Index

- ACF, *see* autocorrelation function
- adaptive Metropolis, 91
  - for marginal sampling, 93
- adaptive proposal, 91
- adjoint method, 108–109
- AM, *see* adaptive Metropolis
- anisotropic
  - diffusion, 60
  - increments, 60
  - prior, 60, 63
  - total variation, 66
- array, 35
- autocorrelation function, 87
  
- Bayes' law, 54, 61, 82, 106, 119
- BCCB, *see* block circulant with circulant blocks
- bias, 10, 19
- biharmonic operator, 70, 71
- block circulant with circulant blocks, 37, 39, 57, 63
  - diagonalize, 39
- block Toeplitz with Toeplitz blocks, 37
- boundary conditions, 3
  - data driven, 28, 40
  - Neumann, 28, 42
  - periodic, 26, 37
  - zero, 3, 34
- BTTB, *see* block Toeplitz with Toeplitz blocks
- burn-in, 84
  
- cdf, *see* cumulative distribution function
- Cholesky factorization, 78–80, 93
- circulant matrix, 26
  - diagonalize, 27, 30
- computed tomography, 42
- conditional autoregression, 54
  
- conjugate gradient iteration
  - preconditioned, 45
  - sampling from the posterior, 81
  - within hierarchical Gibbs, 97
- conjugate prior/hyper-prior, 83
- convolution, 2, 26, 33
- costate equation, 109
- covariance
  - empirical or sample, 91
- covariance matrix, 6
- cross validation, *see* generalized cross validation
- CT, *see* computed tomography
- cumulative distribution function, 117
  
- data-driven boundary conditions, 28, 40
- data models
  - linear, additive Gaussian, 1
  - nonlinear, additive Gaussian, 105
- deblurring, *see* deconvolution
- deconvolution, 2, 33
- degeneracy in hierarchical Gibbs, 88
- detailed balance, 90
- DFT, *see* discrete Fourier transform
- diag operator, 16
- diagonalize
  - block circulant with circulant blocks, 39
  - circulant, 27
- difference matrix, 5
  - Neumann boundary condition, 58
  - periodic boundary condition, 63
- discrepancy principle, 22–23, 37, 46
  
- discrete Fourier transform, 27, 39
- discrete Picard condition, 13
- discretization
  - computed tomography, 43
  - convolution, 3, 26, 34
  - kernel reconstruction, 5
  - negative-Laplacian, 72–73
- DP, *see* discrepancy principle
  
- edge-preserving prior, 63
- eigenvalue decomposition, 78
- empirical covariance, 91
- empirical quantile, 80
- essential sample size, 87
- expected value, 6
  
- fast Fourier transform, *see* discrete Fourier transform
- filter factors, 16
  - Landweber iterations, 18
  - Tikhonov, 17
  - TSVD, *see* truncated singular value decomposition
- filtered SVD, 16
- finite difference discretization of negative-Laplacian, 72–73
- full conditionals, 54, 83, 119
- full posterior density function, 83, 119
  
- gamma distribution, 84
- Gauss–Newton
  - Hessian, 107, 108
  - method, 107
- Gaussian Markov random field, 54–61
  - anisotropic, 59–61, 63
  - edge-preserving, 63
  - higher-order, 70
  - iid increments, 58

- intrinsic, 58
- Neumann boundary condition, 58
- periodic boundary condition, 57
- sampling from, 78
- zero boundary condition, 56
- Gaussian posterior density
  - sampling from, 79–81
- Gaussian random vector, 6
  - nonlinear transformation of, 110
- GCV, *see* generalized cross validation
- generalized cross validation, 21–22, 40, 46, 62
- generalized inverse, 8
- Geweke test, 84
- Gibbs sampler
  - hierarchical, *see* hierarchical Gibbs sampler
  - partially collapsed, 95
  - two-stage, 83
- GMRF, *see* Gaussian Markov random field
- gradient, 107, 108
- gradient scan Gibbs, 98
  
- Hessian, 68, 107, 108
- hierarchical Gibbs sampler, 83
  - degeneracy, 88
  - with embedded PCG, 97
  - with embedded RTO-MH, 118
- hierarchical modeling, 81, 118
- hyper-parameters, 120
- hyper-prior probability density, 82
  
- IACT, *see* integrated autocorrelation time
- iid, *see* independently and identically distributed, 65
- ill-posedness, 10
  - inequality, 7, 10, 11, 106
- increments, 59
  - anisotropic, 60
  - first order, 58
  - iid Gaussian, 58
  - Laplace, 65
  - second order, 70
- independence proposal, 91
- independently and identically distributed, 1
- infinite-dimensional limit
  - hierarchical Gibbs degeneracy, 88
  - well-defined posterior density, 68
- integrated autocorrelation time, 87
- invariant density, 90
- inverse problems, 2
  - test cases, *see* test cases
- iterative regularization, 18
  
- Jacobian, 68, 107, 108, 110, 112, 115
  
- Kaczmarz’s method, 50–51
- kernel, 2, 33
  - Gaussian, 3, 34
  - separable, 34
- kernel reconstruction test case, 4
  - with positivity constraints, 116
- Kronecker product, 35
  
- L-curve, 23–25, 46
- lagged-diffusivity fixed point iteration, 67
- Landweber iteration, 18
- Laplace prior, 65
  - transformation, 117
- Laplacian, negative, *see* negative-Laplacian
- least squares estimator
  - linear, 6
  - nonlinear, 105
  - normal equations, 6
  - statistical properties, 7, 8
  - SVD form, 8
- Levenburg–Marquardt, 106–108
- lexicographical ordering, 35, 63
- likelihood function, 6, 54, 82, 105
- LM, *see* Levenburg–Marquardt
- log-normal prior, 116
  
- MAP estimator, *see* maximum a posteriori estimator
- marginal density, 84, 89
- Markov chain diagnostics, *see* MCMC chain diagnostics
- Markov chain Monte Carlo method, 77
- matrix
  - block circulant with circulant blocks, 37, 39, 57, 63
  - block Toeplitz with Toeplitz blocks, 37
- circulant, 26
- covariance, 6
- difference, 5, 58, 63
- precision, 55
- symmetric positive definite, 56, 78
- Toeplitz, 3, 34
- maximum a posteriori estimator, 54, 61, 106
- maximum likelihood estimator, 6, 105
- MCE, *see* Monte Carlo error
- MCMC chain diagnostics
  - autocorrelation function, 87
  - burn-in, 84
  - equilibrium, 85
  - essential sample size, 87
  - Geweke test, 84
  - integrated autocorrelation time, 87
- MCMC method, *see* Markov chain Monte Carlo method
- mean, 6
- mean squared error, 9, 19
- Metropolis–Hastings, 90–91
  - within hierarchical Gibbs, 118
  - with RTO proposal, 111
- MH, *see* Metropolis–Hastings
- midpoint quadrature, 3, 5, 12, 26, 33, 38
- Monte Carlo error, 86
- MSE, *see* mean squared error
  
- negative-Laplacian
  - finite difference discretization, 72–73
  - Neumann boundary conditions, 59
  - periodic boundary conditions, 57
  - zero boundary conditions, 56
- neighborhood system, 55
- Neumann boundary conditions
  - deconvolution, 28
  - GMRF, 58
  - negative-Laplacian, 59
- Newton’s method, 67, 106
- nonlinear least squares, 105, 106
- normal equations, 6
  
- one-block algorithm, 93
- optimization problems
  - MAP estimator, linear, 67

- MAP estimator, nonlinear, 106
- randomize-then-optimize, 111
- outer product form
  - pseudoinverse, 8
  - SVD, 8
- partially collapsed Gibbs, 95
- PCG, *see* preconditioned conjugate gradient
- periodic boundary conditions
  - convolution, 26, 37
  - GMRF, 57
  - negative-Laplacian, 57
- point spread function, *see* kernel
- Poisson equation inverse problem, 113, 120
- positivity constraints, 116
- posterior density function, 54, 106
  - hierarchical, full, 83, 119
  - sampling from Gaussian, 79–81
- precision
  - matrix, 55
  - parameter, 82
- preconditioned conjugate gradient, *see* conjugate gradient iteration
- preconditioner, 45, 65
- prior
  - improper, 58
- prior probability density
  - edge-preserving, 63
  - Gaussian Markov random field, 54–61, 82, 106
  - iid Gaussian, 54
  - Laplace (total variation), 65
  - log-normal, 116
  - smoothness, 55
  - uniform, 92
  - Whittle–Matérn class, 71
- proposal
  - adaptive, 91
  - independence, 91, 111
  - symmetric, 91
- proposal density, 90
- pseudoinverse, 8
- QR-factorization, 110
- quasi-Newton method, 68
- Radon transform, 43
  - discrete, 43
- randomize-then-optimize, 109–112
- Metropolis–Hastings, 111
- MH within hierarchical Gibbs, 120
- optimization problem, 111
- probability density, 110
- randomized trace estimation, 46
- regression, 1
  - linear, 1
  - principle component, 16
  - ridge, 17
- regularization parameter, 16
- regularization parameter selection methods
  - discrepancy principle, 22–23
  - generalized cross validation, 21–22
  - L-curve, 23–25
  - unbiased predictive risk estimator, 20–21
- regularized solution
  - edge-preserving, 63
  - filtered SVD, 16
  - Landweber iteration, 18
  - statistical properties, 19
  - total variation, 66
  - TSVD, 16
- residual, 107
- ridge regression, 17
- RTO, *see* randomize-then-optimize
- sampling from
  - full posterior density, 83, 95
  - Gaussian Markov random field, 78
  - Gaussian posterior density, 79–81
  - marginal density, 93
  - posterior with Laplace prior, 117
  - posterior with positivity constraints, 116
- semiconvergence, 18
- separable kernel, 34
- signal-to-noise ratio, 4
- singular value decomposition, 7
  - filtered, 16
  - properties, 13
  - truncated, 16
- singular values, 8
- singular vectors, 8
- SNR, *see* signal-to-noise ratio
- SPD, *see* symmetric positive definite
- spectral filtering, 16
- statistical properties
  - least squares solution, 8
  - regularized solutions, 19
- stochastic
  - linear system, 81
  - optimization problem, 111
- SVD, *see* singular value decomposition
- symmetric proposal, 91
- test cases
  - computed tomography, 42
  - deconvolution, 2, 33
  - kernel reconstruction, 5
  - Laplace prior, 117
  - Poisson equation, 113
  - positivity constraints, 116
  - two-parameter, 11, 91, 112, 121
- Tikhonov
  - pseudonormal equations, 17, 42, 44
  - regularization, 17
  - regularized solution, 17, 27, 37, 40, 54
- Toeplitz matrix, 3, 12, 25, 34
- total variance, 9, 19
- total variation, 66
- trace class operator, 69
- trace lemma, 20
- transformation
  - of a Gaussian random vector, 110
  - of the Laplace prior, 117
- transition kernel, 90
- truncated singular value decomposition, 16–17
- unbiased predictive risk estimator, 20–21, 37, 46
- underdetermined, 3, 28
- uniform
  - distribution, 91
  - prior, 92
- UPRE, *see* unbiased predictive risk estimator
- vec operator, 35
- wavelet transform, Haar, 118
- Whittle–Matérn class of priors, 71
- zero boundary conditions
  - convolution, 3, 34
  - GMRF, 56
  - negative-Laplacian, 56